

# Economic evaluation of family-based therapies for children and adolescents: what do we know?

Sandy Tubeuf<sup>1</sup>, Sophie Guthmuller<sup>2</sup>

<sup>1</sup>Academic Unit of Health Economics, Leeds Institute of Health Sciences, University of Leeds, Leeds - UK

<sup>2</sup>European Commission, Joint Research Centre, European Commission, Joint Research Centre, Ispra (Varese) - Italy

## ABSTRACT

**Purpose:** This paper provides a review of the evidence regarding the economic evaluation of family therapy and family-based therapies when delivered to children and adolescents.

**Methods:** Eight databases were searched for published studies and grey literature. We included studies where both costs and effectiveness were evaluated. Due to the substantial differences in interventions, setting, disease area, and sample, we conducted a narrative summary.

**Results:** Sixteen studies were included in the review. Most studies undertook a cost-effectiveness analysis (CEA;  $n = 14$ ) and among those, 1 study additionally carried out a cost-utility analysis. Four studies evaluated the cost-benefit analysis (CBA) of family based intervention, two studies only carried out a CBA while the 2 other studies included both CBA and CEA. The review showed the broad types of family-based interventions and the range of psychological disorders and addictions for which they can be used. Family-based interventions focusing on children and adolescents' health disorders were found to be cost beneficial for all the studies that used CBA, and to be cost effective in two-thirds of the studies that used CEA.

**Conclusions:** Recommendations resulting from the review are that future research should include health care costs beyond costs to the health care sector related to delivering the therapy and the spillover effects to the rest of the family, and should consider cost and effectiveness over a longer time horizon.

**Keywords:** Adolescents, Children, Economic evaluation, Family, Family therapy

## Introduction

Economic evaluation is a key component of health care decision making in the appraisal of new interventions, services, and technologies. It is particularly relevant in England where the National Institute for Health and Care Excellence (NICE) recommends which drugs and treatments the National Health Service (NHS) should fund. They base their guidelines on a methodological "reference case" that favors one type of economic evaluation, namely cost-utility analysis (CUA) with fully incremental analysis (1). This paper reviews the evidence on the economic evaluation of family-based therapy as an intervention for children and adolescents. There are various forms of family-based therapy; it commonly focuses

on relationships within a family and is not solely delivered to the children and adolescents but it also involves their family members. Family-based therapy aims at maximizing cohesion, attachment, adaptability, support, and warmth while reducing maltreatment and scapegoating, and moderating parental control (2). Family-based therapy is manual based and delivered by trained therapists. Several studies have already shown the effectiveness of family therapy and family-based therapy against other types of psychological support or treatment for young people or children for various mental health conditions, including reducing overweight, obesity, self-harm, and suicidal behavior (3-6). A number of other studies have investigated the cost of delivering family therapy using administrative records within a large health care organization, insurance claims, or the Medicaid system, and found it less expensive, which led to fewer repeated episodes of care (7-11). However, the joint evidence of the cost and the effectiveness of family therapies appears to be modest despite the high relevance for policy to establish the value for money that family-based therapies and – more generally – any psychotherapeutic intervention could represent.

Reviews on economic evaluation with a larger scope, including any types of mental health interventions or services for children and adolescents, have already been undertaken (12-16). The first contribution to the area was made by Knapp (12), who identified a handful of literature references and

**Accepted:** August 4, 2017

**Published online:** September 6, 2017

### Corresponding author:

Dr. Sandy Tubeuf  
Academic Unit of Health Economics  
Leeds Institute of Health Sciences  
University of Leeds  
Level 11, Worsley Building  
Leeds LS2 9NL, UK  
s.tubeuf@leeds.ac.uk

made a compelling argument for increasing the use of economic evaluation in children and adolescents' mental health decisions. He demonstrated how commissioners, clinicians, and researchers relied on economic analysis to obtain the best value from resources allocated to the planning and provision of services for children and adolescents with mental health problems. The general conclusion of the reviews of the past 20 years is that the number of studies where both costs and health outcomes of mental health interventions and services for children and adolescents are assessed has increased, but still remains limited, particularly compared to mental health interventions in the adult population.

Our review differs from previous reviews in 2 ways: it focuses on 1 type of intervention: family therapy or family-based therapy for children and adolescents, and it is not limited to a specific type of mental health or health condition.

We describe the methods we used for the identification of relevant studies, present the results of the literature review, and then discuss these results. We conclude with an exploration of further avenues of research.

## Methods

### Search strategy

The literature search aimed to identify studies on economic evaluations of family-based intervention therapy for children and adolescents. We searched the following databases in February 2015 from their inception dates to the present: CINAHL (EBSCO), Cost Effectiveness Analysis Registry (CEVR, Tufts Medical Centre), Embase Classic, Embase (Ovid), IDEAS (RePEc), NHS Economic Evaluation Database, Ovid MEDLINE (R), and PsycINFO (Ovid).

The searches were designed to identify studies on the cost and the effectiveness of family-based intervention for young persons by combining the search concept "children" or "adolescent," and "family therapy" or "family intervention," with the concepts "economic evaluation" or "cost effectiveness." A full detail of the search strategy for Medline database is presented in Appendix A (available online at [www.grhta.com](http://www.grhta.com)) and full search strategies for all databases are available from the authors upon request. Bibliographies of included studies were also searched for relevant studies.

### Inclusion criteria

To be eligible for inclusion, the studies had to evaluate both costs and effectiveness. We used the economic evaluation CHEERS checklist developed by Husereau et al (17) as a guide for best practice when conducting economic evaluations, and we only selected studies for full review where the 2 components of an economic evaluation were measured: costs and health-related effectiveness. Evaluations of family-based interventions focusing on cost only or on outcomes only were excluded. In addition, we excluded studies that reported qualitative data and literature reviews; studies that lacked quantitative data; studies in developing countries; and studies in which the language was not English. Also, studies that evaluated family-based intervention for adults, or studies where the impact of parenting interventions was measured on parents

only were excluded, as the main focus of the review was to analyze the impact on children or young people. We did not apply any specific criteria for the family-based therapy and included any therapy where the family was involved. Similarly, we did not specify inclusion criteria for the disease areas, assuming that family-based therapies were likely to focus on mental health and behavioral disorders. Finally, we did not set any publication date limit, and studies were searched spanning almost 70 years (1946-2015).

### Data collection and analysis

To select studies for the review, titles and abstracts were screened by 1 researcher using the above inclusion criteria; this process was managed using Endnote Software. When the decision to include or exclude a study was not straightforward, a second reviewer was consulted to resolve uncertainty. The detailed process of article selection was presented in a PRISMA chart (18). Once the included studies were identified, data were extracted using a form developed for this purpose. For each study, we extracted the following: cost perspective, type of economic evaluation, study population, country, outcome measures and length of follow-up, type of family based intervention, and sensitivity analysis. Due to the substantial differences in interventions, settings, disease areas, and samples, a meta-analysis was not conducted. Instead, a narrative summary of the included studies is presented below.

## Results

The literature search yielded 1,322 potentially relevant studies. After the removal of duplicates, 920 references were identified for screening. In addition, 3 studies were identified scanning from a related review. In total, 923 titles and abstracts were screened of which 886 studies were excluded and 37 possibly relevant studies were retrieved for full-text assessment. After full-text review, 16 studies were included in the analysis and 21 studies were excluded. Studies were excluded because: 6 only focused on outcomes; 9 evaluated a parenting program; 4 were a family-based intervention for adults; 1 did not include family therapy as an intervention; and 1 was not a cost-effectiveness analysis (CEA). A PRISMA chart describing the inclusion and exclusion process is shown in Figure 1.

### Description of included studies

Of the 16 studies included in the review, 9 were from the USA, 3 from the UK, and 1 each from the Netherlands, Canada, Sweden, and Finland (Tab. I). The majority of the studies were published during the last 10 years (9 studies from 2007 to 2014), 4 in the first half of 2000, and 3 before 2000 (1988, 1989, 1999).

The studied populations comprised children and/or adolescents aged between 1 and 20 years old. Family-based interventions were assessed in four main disease areas; (i) eating disorders, such as obesity or anorexia ( $n = 4$ ); (ii) mental health or psychosomatic disorders, including anxiety, deliberate poisoning, or asthma ( $n = 5$ ); (iii) substance use disorders




Fig. 1 - Flow diagram of excluded and included studies.

TABLE I - Methodology of included studies

Study reference	Disease area	Country	Study population	Sample size	Study design	Data sources	Follow-up
Byford et al (32)	Deliberate poisoning	UK	Under 17	n = 162	CEA	RCT	6 months
Goldfield et al (25)	Obesity	USA	Aged 8-12	n = 24	CEA	RCT	12 months
French et al (28)	Substance use disorder	USA	Aged 13-17	n = 114	CEA	RCT	7 months
Bodden et al (19)	Anxiety	Netherlands	Aged 8-18	n = 128	CEA/CUA	RCT	12 months
Flores et al (34)	Asthma	USA	Aged 2-18	n = 220	CEA	RCT	12 months
Byford et al (26)	Anorexia	UK	Aged 12-18	n = 135	CEA	RCT	24 months
Dennis et al (29)	Cannabis use disorder	USA	Aged 15-16	n = 600	CEA	2 RCTs	12 months
Kalavainen et al (27)	Obesity	Finland	Aged 7-9	n = 70	CEA	RCT	12 months
Dopp et al (20)	Juvenile offender	USA	Aged 12-17	n = 176	CBA	RCT + tracking	25 years
Klietz et al (21)	Juvenile offender	USA	Aged 12-17	n = 176	CBA	RCT + tracking	13.7 years
Schmidt et al (31)	Eating disorder	UK	Aged 13-20	n = 85	CEA	RCT	12 months
Gustafsson & Svedin (22)	Psychosomatic disorder	Sweden	Aged 1-16	n = 42	CEA/CBA	Retrospective study	5 years
Moore & Crane (33)	Relational disorder	USA	Aged 1-18	n = 960	CEA	Retrospective study	4 years
Spoth et al (30)	Alcohol use disorder	USA	Sixth graders	n = 478	CEA	RCT	12 months
Dydyk et al (35)	Symptomatic behavior	Canada	Aged 3-14	n = 42	CEA	RCT	12 months
Crane et al (9)	Conduct disorder	USA	Aged 5-18	n = 3753	CBA	Retrospective study	30 months

CBA = cost-benefit analysis; CEA = cost-effectiveness analysis; CUA = cost-utility analysis; RCT = randomized controlled trial.



(alcohol, cannabis) ( $n = 3$ ); and (iv) conduct disorders, including juvenile offenders ( $n = 4$ ). Most of the analyses ( $n = 11$ ) used data from randomized control trials. They included, on average, 187 (standard deviation [SD]: 184.4) participants, and the follow-up period ranged from 6 months to 24 months with most trial studies using a 12-month follow-up ( $n = 8/11$ ). Three studies were based on retrospective cohort studies and included a larger number of participants (on average 1,585 [SD: 1932]) that were followed between 30 months and 5 years. Finally, the 2 juvenile offender studies relied on a sample of participants from a trial, which were then followed on a longer term for a cost-benefit analysis (CBA). They each included 176 participants and the duration for the follow-up was 13.7 years and 25 years, respectively.

Most studies undertook a CEA ( $n = 14$ ) and among those, 1 study additionally carried out a CUA (19). Four studies evaluated the CBA of family-based intervention, 2 studies only carried out a CBA (20, 21), while the 2 other studies included both CBA and CEA (9, 22).

### *Design of interventions*

The studies included in the review showed a variety of designs of family-based interventions including family therapy ( $n = 5$ ), multi-systemic therapy ( $n = 2$ ), functional family therapy, multidimensional family therapy, intensive family therapy, joint family therapy and cognitive behavioral therapy (CBT), family CBT, and other family-based programs (Tab. II). There were 3 main characteristics that appeared to distinguish between family therapies in general. First, therapies could be delivered either in group (several families in a session with 1 therapist) or in individual sessions (1 family with 1 therapist) (23). Second, the intervention could be based at home (therapist's visits), at the hospital, or in a community practice. Finally, sessions could be delivered to the whole family (parent + child) or in separate session (parent only, or child only). Separate session for parents or caregivers usually took the form of parental counselling with a therapist (24).

In more than half of the studies, family-based interventions combined treatment sessions with the whole family, sessions with only 1 parent or caregiver, and additional sessions with the treated child only. Group sessions, where parents and children were seen separately, were most often used in the treatment of eating disorders (25-27). On the other hand, family therapies designed to treat mental health disorders or substance use disorders usually comprised family sessions with a therapist.

Comparison groups for the study included treatment as usual ( $n = 7$ ), individual therapy ( $n = 5$ ), individual CBT ( $n = 3$ ), group CBT ( $n = 1$ ), and a mix of both individual and group therapy. Several studies evaluated more than 1 family-based intervention and presented a study design with more than 2 treatment arms (9, 26, 28-30).

### *Outcome measures*

Figure 2 presents the frequency of the outcomes and the cost measurements used across the 16 studies, and Table I in Appendix B (available online at [www.grhta.com](http://www.grhta.com)) provides an extended description of those. In the majority (13 out of 16)

of the included studies, effectiveness was measured using health outcome measures. More specifically, the effectiveness of family interventions in eating disorder conditions was assessed using weight loss, such as the reduction in body mass index units (25, 27) and a change in the Morgan-Russell Average Outcome Scale (MRAOS) for patients with anorexia (26), or the abstinence rate from bingeing or vomiting at the end of the follow-up period (31). In the area of substance use disorders, the effectiveness of family-based interventions was evaluated as a change in the number of days of abstinence from cannabis use (29), the abstinence from marijuana use (28), or the number of case of alcohol use disorder prevented (30). For mental health disorders, health outcomes were measured as a change in the suicidal ideation scale and hopelessness (32), the number of anxiety-free days (19), or the return rate to therapy (33). Finally, the study on asthma used the reduction in asthma symptoms as a health outcome (34). Some studies used health care use and the reduction in hospital length of stay as measures of effectiveness (22, 35).

CBA measured the effectiveness in monetary value and was used in 4 out of 16 studies. In 2 studies, it represented avoidance of crime victim expenses related to family interventions for juvenile offenders and savings to the taxpayer (20, 21). The 2 other studies combining both CEA and CBA considered effectiveness as an overall reduction in health care expenses induced by family therapy for children with conduct disorders (9, 22).

Utility values were measured in 1 study only (19); quality-adjusted life years (QALYs) were generated from the generic health-related quality of life instrument EQ-5D (36) to evaluate the effectiveness of family therapy compared to individual therapy in the treatment of anxiety, in addition to the proportion of anxiety-free days.

As family therapy involves parents, care givers, brothers, sisters, or other close relatives and friends, 4 papers had interestingly looked at the impact of the treatment on relatives participating in the therapy in addition to the impact of the therapy on the adolescent/child patient. In 1 study (32), parents completed the General Health Questionnaire along with a treatment satisfaction questionnaire. They found that parents randomized in the family-based intervention were more satisfied than parents of the controls for children with deliberate self-poisoning at 2 months follow-up. Another study (25) looked at the impact of family therapy on the weight loss of parents along with the reduction of weight of children. Similarly, the asthma study (34) investigated the effectiveness of the interventions on the quality of life of care givers using the Pediatric Asthma Caregiver's Quality of Life Questionnaire (PACQLQ). Finally, only 1 study assessed the effectiveness of the intervention on the whole family. Bodden et al (19) collected EQ-5D scores of parents and children, and calculated family-based QALYs to evaluate the effectiveness of the intervention for clinically anxious children.

### *Cost measures*

Economic evaluation studies require that cost components are specified precisely and presented in a disaggregated form. All 16 studies included the direct cost related to the

**TABLE II** - Design of the family-based therapy

Study reference	Disease area	Design of the family-based therapy		Design of the comparator	
		Name of the intervention	Description	Name of the intervention	Description
Byford et al (32)	Deliberate poisoning	Home-based social work (Home FT)	Home FT in addition to the routine care. One assessment session and four intensive family-centered, home-based intervention sessions with two psychiatric social workers	Routine care or TAU	Out-patient clinic visits with psychiatrists and psychiatric nurses.
Goldfield et al (25)	Obesity	Group treatment	13 sessions with children and parents receiving group treatment separately.	Mix of individual treatment and group therapy (Mixed)	Sessions of 15-20 mins individual sessions with a therapist and 40 mins of group therapy.
French et al (28)	Substance use disorder	FFT	Family therapy	Ind. CBT	CBT
		Joint individual CBT + FFT	Joint integrates treatment combining individual and family therapy	Group CBT	Skills focused psychoeducational group treatment
Bodden et al (19)	Anxiety	Family CBT	12 sessions	Ind. CBT	12 sessions
Flores et al (34)	Asthma	PM	One home visit + monthly meeting at a community learning center with children and family	TAU	Usual pediatric asthma care
Byford et al (26)	Anorexia	Specialist out-patient treatment	Ind. CBT + parental counselling with the participant + dietary therapy and multimodal feedback.	In-patient psychiatric treatment	Multidisciplinary psychiatric approach with the aim of normalizing eating, restoring healthy weight and facilitating psychological cognitive change.
		General out-patient treatment	TAU in community CAMHS with multidisciplinary, individual- and family-based approach, with variable dietetic and pediatric liaison		
Dennis et al (29)	Cannabis use disorder	FSN	MET/CBT12 + 6 parent education group meetings, 4 therapeutic home visits, referral to self-help support group and case management for 12 to 14 weeks	MET/CBT5	5 sessions (2 individual MET and 3 group CBT sessions) during 6 to 7 weeks
		ACRA	10 individual sessions, 4 sessions with the caregivers and limited case management for 12 to 14 weeks	MET/CBT12	7 additional CBT sessions in group format
		MFT	12 to 15 sessions (6 with the adolescent, 3 with the parents and 6 with the whole family) + case management for 12 to 14 weeks		
Kalavainen et al (27)	Obesity	Group program	14 sessions held separately for parents and children	Routine program (TAU)	Two individual appointments for the children
Dopp et al (20)	Juvenile offender	MST	MST delivered to youths and their caregivers in home, school or neighborhood settings for 4 to 6 months	IT	Usual community outpatient treatment for juvenile offenders
Klietz et al (21)	Juvenile offender	MST	MST delivered to youths and their caregivers in home, school or neighborhood settings for 4 to 6 months	IT	Usual community outpatient treatment for juvenile offenders
Schmidt et al (31)	Eating disorder	FT	FT for anorexia nervosa including family and individual sessions. Up to 13 sessions with close others and 2 individual sessions over a 6-month period	CBT-guided self-care	Manual + 10-weekly sessions

To be continued



TABLE II - Continued

Study reference	Disease area	Design of the family-based therapy		Design of the comparator	
		Name of the intervention	Description	Name of the intervention	Description
Gustafsson & Svedin (22)	Psychosomatic disorder	FT	FT included the whole family or part of it according to the structural family model of Minuchin, Rosman, and Baker. Individual sessions with selected children for 6 months	TAU	Children hospitalized with the same diagnosis
Moore & Crane (33)	Relational disorder	FT or Mixed	FT and Mixed identified by CPT code for parent-child relational problem + a relational CPT code in Cigna	IT	IT identified by current procedural terminology code for parent-child relational problem in Cigna
Spoth et al (30)	Alcohol use disorder	ISFP	Seven sessions with children in all sessions and includes more parent-child interactive activities than does PDFY	TAU	Minimal contact condition
		PDFY program	Five sessions		
Dydyk et al (35)	Symptomatic behavior	IFTU	Multiple impact therapy with families: family sessions, overlapping interviews, subsystem work, individual sessions, assigning of task, use of audio-visual feedback, observation and problem solving in instrumental areas such as bedtime, meals and use of leisure time	TAU	Comparison families
Crane et al (9)	Conduct disorder	In-office FT	FT in office 30-month follow-up	TAU	Youth diagnosed with conduct disorder received a variety of services but not FT
		Behavior management in home FT	FT at home, 30-month follow-up		

ACRA = adolescent community reinforcement approach; CAMHS = children and adolescents mental health services; CBT = cognitive behavioral therapy; CPT = current procedural terminology; FFT = functional family therapy; FT = family-based therapy; FSN = family support network; IFTU = intensive family therapy unit; Ind. CBT = individual CBT; ISFP = Iowa Strengthening Families Program; IT = individual therapy; MET = motivational enhancement treatment; MFT = multidimensional family therapy; Mixed = mixed therapy; MST = multisystemic therapy; PDFY = preparing for drug-free years; PM = parent mentor; TAU = treatment as usual.

delivery of the intervention; however, only 6 studies included additional costs.

One study (9) adopted a health-sector-cost perspective, including all Medicaid billed costs for pharmacy, hospital, and professional services for 30 months. Two studies adopted a broad service-providers' perspective, and included costs related to any use of health, social, and educational services within the public perspective. More specifically, Byford et al (32) considered the costs of using health and social services, including hospital services, primary care, social worker and community care, foster and residential care, counselling by practice nurses, clinical psychologists or school counsellors, as well as the costs of educational welfare officers, psychologists, school doctors, and school nurses over the 6-month study period. Similarly, Byford et al (26) considered costs related to health, social services, education, and the voluntary and private sectors, including hospital contacts, private-sector inpatient stays, community health and social services, schooling, and medications at the 1- and 2-year follow-up points in the study.

The 3 other studies additionally considered costs to the family and adopted a societal perspective on costs (19, 31,

34). Bodden et al (19) included direct health care costs (e.g., contact with the GP, psychologist, psychiatrist, and the medication), direct non-health-care costs (e.g., paid housekeeper, day nursery, informal care), indirect costs (paid work, unpaid work, loss of leisure time, household work, school absence) and out-of-pocket costs related to health care use. The cost diaries used in the study recorded the resources used by all family members (child, sibling, mother, or father) as well as the reason for the resource use at the 3- and 6-month follow-up points; their analysis then only included family resource use related to the child's anxiety. Flores et al (34) estimated the total cost of the intervention including the program costs (personnel involved, payments to parents mentors, training sessions, monthly meetings, bus fares), direct medical costs (charges to emergency department visits and hospitalizations), and indirect income costs of missed work days for the employed caregivers. Finally, Schmidt et al (31) included adolescents' public sector costs, such as education (e.g., home tuition, individual help in class, contact with the school nurse, educational psychologist, educational welfare officer, meeting with tutors, any educational support), health and social care service costs




Fig. 2 - Types of outcomes and costs studied in the literature.

for the child (e.g., hospital care, primary care, A&E care; specialist services, e.g., dietician, psychiatrist, and child development center, social worker), and costs for family members (GP, outpatient appointments, and psychiatrist or psychologist). In addition, they included costs to the family and the patient as a consequence of bulimia nervosa, such as the cost of lost employment and families’ and patients’ out-of-pocket expenses. The costs were collected at baseline, and at 6- and 12-month assessments.

**Is family therapy cost effective?**

The results of the economic evaluation for each study are summarized in Table III. At the end of the follow-up period, of the 16 included studies, 11 concluded that family-based therapy was clearly cost beneficial and cost effective compared with the different comparators, and for various types of health outcomes, while another 2 studies found it cost effective under specific conditions. Byford et al (32) found no difference in cost or in effectiveness on the whole sample, but found that family therapy improved suicidal ideation for a subgroup of patients with major depression. Similarly the paper by Byford et al (26) showed no significant difference in the cost-effectiveness of family therapy compared to routine care in hospital, but the family-based outpatient treatment had the highest probability of being cost effective when uncertainty was taken into account. Family-based therapy was found to be cost beneficial for studies on juvenile offenders (20, 21) and cost effective for conduct disorders (9). It was also found to be cost effective for most of the studies on mental health, psychosomatic, and relational disorders. We note

that family therapy was found both cheaper and more effective than treatment as usual for patients with asthma (34) and patient with psychosomatic disorders (22), and it was found cheaper and with equal effectiveness than the comparator for adolescents with symptomatic behaviours (35) and relational disorders (33). An exception to the cost effectiveness of family-based therapy for mental health disorders is the study on clinically anxious children (19), which showed that family CBT compared with individual CBT was of equal cost and effectiveness, but concluded that individual CBT had the highest probability of being cost effective in both cost per anxiety-free child and cost per QALY when modelling the CEA.

While family-based therapy is found to be cost effective for children with obesity (25, 27) and anorexia (26), it was found to be more expensive and equally effective as self-guided CBT for adolescents with bulimia nervosa (31). Schmidt et al (31) concluded that family therapy is more costly and equally effective than individual self-guided CBT in the treatment of eating disorders. Most of the studies on substance use – except the study on alcohol – did not give evidence that family-based therapies were cost beneficial or cost effective when compared to another intervention, and this was mainly due to family-based therapies being more expensive than the comparator, and of equal or lower effectiveness. Dennis et al (29) found that the Family Support Network was more costly and less effective in the treatment of a cannabis use disorder. French et al (28) found no difference in effectiveness between treatments including family-based, individual, and group cognitive behavioral approaches. However, individual group-based therapy are the least costly compared to functional family therapy.



TABLE III - Is the family-based therapy cost effective?

Study reference	Disease area	Outcome measure	Family-based therapy	Comparator (C)	Cost	Effectiveness	CE result <sup>1</sup>
Byford et al (32)	Deliberate poisoning	Suicidal ideation and hopelessness scale	Home FT	TAU	Home FT = TAU	Home FT = TAU	FT (subgroup)
Goldfield et al (25)	Obesity	Reduction in BMI & % overweight	Group FT	Mixed therapy	Group FT < Mixed	Group FT = Mixed	FT
French et al (28)	Substance use disorder	% of days of marijuana use & delinquency score	FFT or joint FFT + CBT	CBT or group	Joint>FFT>CBT>Group	Joint = FFT = CBT = Group	C (group)
Bodden et al (19)	Anxiety	Anxiety free rate and QALY	Family CBT	Ind. CBT	Family CBT = Ind. CBT	Family CBT = Ind. CBT	Proba C
Flores et al (34)	Asthma	Reduction in asthma symptoms	PM	TAU	PM<TAU	PM>TAU	FT
Byford et al (26)	Anorexia	Change in MRAOS	Specialist or General out-patient	In-patient	G. Out- = Spe. Out- = In-	G. Out- = Spe. Out- = In-	Proba FT
Dennis et al (29)	Cannabis use disorder	Days of abstinence	FSNM	METS5 or MET12	FSNM>METS5=MET12	FSNM<METS5=MET12	C
	Cannabis use disorder	Days of abstinence	ACRA or MDFT	METS5	ACRA<METS5<MDFT	MDFT<METS5<ACRA	FT
Kalavainen et al (27)	Obesity	1% weight for height decrease & change in BMI-SDS unit score	Group FT	TAU	Group FT>TAU	Group FT>TAU	FT
Dopp et al (20)	Juvenile offender	Taxpayer and crime victim expenses avoided	Multisystemic therapy	Ind. therapy	MST>IT	MST>IT	FT
Klitz et al (21)	Juvenile offender	Taxpayer and crime victim expenses avoided	Multisystemic therapy	Ind. therapy	MST>IT	MST>IT	FT
Schmidt et al (31)	Eating disorder	Abstinence rate from bingeing or vomiting	FT	CBT self-guided	FT>CBT self-guided	FT = CBT self-guided	C
Gustafsson & Svedin (22)	Psychosomatic disorder	Reduction of number of days at hospital	FT	TAU	FT<TAU	FT>TAU	FT
Moore & Crane (33)	Relational disorder	Return rate to therapy	FT or mixed therapy	Ind. therapy	Mixed>FT = IT	FT = IT = MT	FT
Spoth et al (30)	Alcohol use disorder	Cost per case prevented per 100 families treated	ISFP or PDFY	TAU	ISFP>PDFY>TAU	ISFP>PDFY>TAU	FT
Dydyk et al (35)	Symptomatic behaviour	Reduction of length of stay in residential care	IFTU	TAU	IFTU<TAU	IFTU>TAU	FT
Crane et al (9)	Conduct disorder	Costs of health care	In-office or in-home FT	TAU	Not provided	In-home > in-office>TAU	FT

<sup>1</sup> Cost effectiveness result at the end of the last follow-up period.

ACRA = adolescent community reinforcement approach; BMI = body mass index; BMI-SDS = BMI standard deviation score; CBT = cognitive behavioral therapy; FFT = functional family therapy; FSNM = family support network meetings; FT = family-based therapy; Ind. CBT = individual CBT; ISFP = Iowa Strengthening Families Program; MDFT = multidimensional family therapy; MRAOS = Morgan-Russell Average Outcome Scale; PDFY = preparing for drug-free years; TAU = treatment as usual; PM = parent mentor; QALY = quality-adjusted life year.



## Discussion

The results from the literature review show diversity of 2 kinds. First, there is diversity in the types of family-based intervention being evaluated as previously stated by Pote et al (37). Second, family-based intervention is used to treat a broad range of psychological disorders and addictions among children and adolescents. Based on the included studies, we found that family-based interventions focusing on children and adolescents' health disorders are cost beneficial for the 3 studies that used CBA, and are cost effective in two-thirds of the studies that used CEA. If we focus on the best practice guidelines for conducting economic evaluations (1, 38), this review shows that the quality of economic evaluations of family-based interventions for young people and children is limited. We develop this further in this section.

In the majority of the papers, the cost analysis only included the cost to the health care sector to deliver the intervention. Few papers included health resource use consecutive to the intervention (e.g., medications, GP visits, etc.) over a key time horizon that is long enough to reflect all the important differences in costs between the treatments being compared. Only 3 papers considered the costs from other public sectors, such as education, justice, or social services, despite 1 study underlying that family-based interventions have an impact on the resource use of children and adolescents in several public sectors, which are important to take into account in economic evaluations (32).

Regarding consequences and, more generally, effectiveness measurement, the NICE reference case recommends the use of a health outcome that not only is reliable for the comparison between competitive interventions, but also is appropriate for measuring the marginal impact of the health care decisions the policy maker will make for a broad range of treatments within the same budget. In this context, health effects should be expressed in terms of QALYs, which is currently considered to be the most appropriate generic measure of health benefit that reflects both mortality and health-related quality of life (1). While NICE states that EQ-5D is the preferred measure of health-related quality of life in adults, there is no consensus on the best health-related quality of life instrument to use in children and young people (39). Recent reviews of literature have identified several preference-based instruments being used to assess children's utility, including the young people modified EQ-5D, the Child Health Utility 9 dimensions (CHU-9D), the Health Utility Index II (HUI II) or the proxy-respondent EQ-5D (40, 41). In our review, only 1 study undertook a CUA using QALYs as the outcome and found individual CBT to be more cost effective than family CBT therapy (19). The paucity of utility data in economic evaluation of children interventions has already been highlighted (40) and this raises an important issue to inform resource allocation decisions across different treatments and health conditions.

With family-based therapies, there is potential to account for additional health benefits for relatives and carers in the economic evaluation. Several studies in the review collected parents or carers' outcomes and used them as additional outcomes of interest in a cost-effectiveness analysis (25, 32, 34), and 1 study combined both children's and parents' outcomes (19). The combination of benefits beyond the patient is not

specifically stated in the NICE reference case, and relies on the strong assumption that QALYs can be aggregated across individuals as a simple additive aggregation. Whilst this has been done in prior studies on child health (42), and would be consistent with other health economic research showing benefits to other family members (43), such considerations require a theoretical justification and a discussion on the interdependence between the utility functions of children and adolescents and of the parents, as well as a consensus on how best to measure utility in children.

In an economic evaluation, the costs and benefits require an optimal length of follow-up or a time horizon that is long enough to reflect substantial differences in costs or consequences between the treatments being compared. If the analysis is undertaken within a trial follow-up, then the economic evaluation will be restricted to that horizon. However, one could easily imagine that therapies, such as family therapies, can process beyond the sessions that are attended with teenagers in the longer term, reflecting on the therapy; therefore such interventions are likely to keep on giving and improving the individual's wellbeing beyond the trial follow-up. Decision analysis modelling is the preferred method in this context; it consists of extrapolating the costs and consequences using probabilities and assumptions based on the data collected in a trial and data extracted from the literature. None of the papers in the review evaluated the long-term economic evaluation of family-based therapies using a decision analysis model. This probably comes from the fact that children and adolescents face such a large number of events over their life, and such a decision analysis model would carry a high level of uncertainty, especially if a lifetime horizon is considered. A mitigating strategy could be to consider a shorter time horizon, such as reaching the age of 25, which is considered to be the age when the adolescent brain is fully developed.

The strengths of this review include conducting a comprehensive search to identify all published and unpublished studies estimating the economic evaluation of family-based therapies for children and adolescents in developed countries. To the best of our knowledge, this is the first review in that area. While we tried to minimize biases during the review, we found that due to the high heterogeneity and diversity of the comparison groups in studies, it was not appropriate to conduct a meta-analysis. In addition, due to limited capacity, we were unable to include non English-language studies. While every effort was made to identify all relevant articles, it is also possible that we may have missed some unpublished grey literature. The aim of this review was to find studies that undertook an economic evaluation of family-based interventions in children and adolescents and it discussed a number of limitations that are specific to the children and adolescents' health context. Therefore, the findings of this review are unlikely to be generalizable to the economic evaluation of family-based therapy in an adult population.

## Conclusion

While a number of studies in North America and Europe have simultaneously considered the cost effectiveness, clinical effectiveness, or health benefits of family-based therapies to treat young people and children with various conditions,

few have conducted an economic evaluation considering costs beyond the delivery of the therapy. As family interventions have been found effective, particularly with children and adolescents (44-46), we suggest that future economic evaluations of family-therapies should include costs related to consecutive health care use over a longer time horizon. Methodological and theoretical concepts currently under development in health economics regarding the best way to account for potential spillover health benefits on families, and to validate utility measurements for the children and adolescent population, will also be key to guide the economic evaluations of family interventions for children and adolescents.

## Acknowledgments

This review research was undertaken as part of the National Institute for Health Research Health Technology Assessment Programme (project number: 07/33). The views expressed in this publication are those of the authors and do not necessarily reflect those of the HTA programme, NIHR, NHS, or the Department of Health. The final revisions of the article have been conducted when Sophie Guthmuller took service at the European Commission, Joint Research Centre. The scientific output expressed does not imply a policy position of the European Commission. Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use which might be made of this publication. The authors would like to thank Judy Wright and Kirstine McDermid for literature searches; and Joachim Marti and two anonymous reviewers for comments on an earlier version of the manuscript.

## Disclosures

Financial support: None.

Conflict of interest: None of the authors has financial interest related to this study to disclose.

## References

- National Institute for Health and Care Excellence. Guide to the methods of technology appraisal. <https://www.nice.org.uk/process/pmg9/chapter/foreword>. Accessed August 21, 2017.
- Fortune S, Cottrell D, Fife S. Family factors associated with adolescent self-harm: a narrative review. *J Fam Ther*. 2016;38(2):226-256.
- Antal H, Buckloh L, Lochrie A, et al. Family-based intervention for overweight youth: Effects on health-related quality of life and measurements of physical health. *Diabetes*. 2010. Conference: 70<sup>th</sup> Scientific Sessions of the American Diabetes Association. Orlando, FL United States.
- De Silva S, Parker A, Purcell R, Callahan P, Liu P, Hetrick S. Mapping the evidence of prevention and intervention studies for suicidal and self-harming behaviors in young people. *Crisis*. 2013;34(4):223-232.
- Vetere A. Evaluation of systemic interventions. In: Brown JM and Campbell EA, eds. *The Cambridge handbook of forensic psychology*. xxiv. New York, NY, US: Cambridge University Press 2010; 813-819.
- Vos RC, Huisman SD, Houdijk EC, Pijl H, Wit JM. The effect of family-based multidisciplinary cognitive behavioral treatment on health-related quality of life in childhood obesity. *Qual Life Res*. 2012;21(9):1587-1594.
- Crane DR. The cost-effectiveness of family therapy: a summary and progress report. *J Fam Ther*. 2008;30(4):399-410.
- Crane DR, Christenson JD, Dobbs SM, et al. Costs of treating depression with individual versus family therapy. *J Marital Fam Ther*. 2013;39(4):457-469.
- Crane DR, Hillin HH, Jakubowski SF. Costs of treating conduct disordered medicaid youth with and without family therapy. *Am J Fam Ther*. 2005;33(5):403-413.
- Law DD, Crane DR. The influence of marital and family therapy on health care utilization in a health-maintenance organization. *J Marital Fam Ther*. 2000;26(3):281-291.
- Morgan TB, Crane DR, Moore AM, Eggett DL. The cost of treating substance use disorders: Individual versus family therapy. *J Fam Ther*. 2013;35(1):2-23.
- Knapp M. Economic evaluations and interventions for children and adolescents with mental health problems. *J Child Psychol Psychiatry*. 1997;38(1):3-25.
- Romeo R, Byford S, Knapp M. Annotation: Economic evaluations of child and adolescent mental health interventions: a systematic review. *J Child Psychol Psychiatry*. 2005;46(9):919-930.
- Zechmeister I, Kilian R, McDavid D. MHEEN group. Is it worth investing in mental health promotion and prevention of mental illness? A systematic review of the evidence from economic evaluations. *BMC Public Health*. 2008;8(1):20.
- Kilian R, et al. Cost-effectiveness analysis and child and adolescent mental health problems. *Int J Ment Health Promot*. 2010;12:45-57.
- Beecham J. Annual research review: Child and adolescent mental health interventions: a review of progress in economic studies across different disorders. *J Child Psychol Psychiatry*. 2014;55(6):714-732.
- Husereau D, Drummond M, Petrou S, et al. CHEERS Task Force. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *Value Health*. 2013;16(2):e1-e5.
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med*. 2009;151(4):W65-94.
- Bodden DH, Dirksen CD, Bögels SM, et al. Costs and cost-effectiveness of family CBT versus individual CBT in clinically anxious children. *Clin Child Psychol Psychiatry*. 2008;13(4):543-564.
- Dopp AR, Borduin CM, Wagner DV, Sawyer AM. The economic impact of multisystemic therapy through midlife: a cost-benefit analysis with serious juvenile offenders and their siblings. *J Consult Clin Psychol*. 2014;82(4):694-705.
- Klietz SJ, Borduin CM, Schaeffer CM. Cost-benefit analysis of multisystemic therapy with serious and violent juvenile offenders. *J Fam Psychol*. 2010;24(5):657-666.
- Gustafsson A, Svedin CG. Cost effectiveness: Family therapy in a pediatric setting. *Fam Syst Med*. 1988;6(2):162-175.
- Wilkins JR III, Crawford JM, Stallones L, et al. Using participant event monitoring in a cohort study of unintentional injuries among children and adolescents. *Am J Public Health*. 2007;97(2):283-290.
- Hightow-Weidman LB, Phillips G II, Outlaw AY, et al. Patterns of HIV disclosure and condom use among HIV-infected young racial/ethnic minority men who have sex with men. *AIDS Behav*. 2013;17(1):360-368.
- Goldfield GS, Epstein LH, Kilanowski CK, Paluch RA, Kogut-Bossler B. Cost-effectiveness of group and mixed family-based treatment for childhood obesity. *Int J Obes Relat Metab Disord*. 2001;25(12):1843-1849.
- Byford S, Barrett B, Roberts C, et al. Economic evaluation of a randomised controlled trial for anorexia nervosa in adolescents. *Br J Psychiatry*. 2007;191(5):436-440.
- Kalavainen M, Karjalainen S, Martikainen J, Korppi M, Linnosmaa I, Nuutinen O. Cost-effectiveness of routine and group programs for treatment of obese children. *Pediatr Int*. 2009;51(5):606-611.
- French MT, Zavala SK, McCollister KE, Waldron HB, Turner CW, Ozechowski TJ. Cost-effectiveness analysis of four interventions



- for adolescents with a substance use disorder. *J Subst Abuse Treat.* 2008;34(3):272-281.
29. Dennis M, Godley SH, Diamond G, et al. The Cannabis Youth Treatment (CYT) Study: main findings from two randomized trials. *J Subst Abuse Treat.* 2004;27(3):197-213.
  30. Spoth RL, Guyll M, Day SX. Universal family-focused interventions in alcohol-use disorder prevention: cost-effectiveness and cost-benefit analyses of two interventions. *J Stud Alcohol.* 2002;63(2):219-228.
  31. Schmidt U, Lee S, Beecham J, et al. A randomized controlled trial of family therapy and cognitive behavior therapy guided self-care for adolescents with bulimia nervosa and related disorders. *Am J Psychiatry.* 2007;164(4):591-598.
  32. Byford S, Harrington R, Torgerson D, et al. Cost-effectiveness analysis of a home-based social work intervention for children and adolescents who have deliberately poisoned themselves. Results of a randomised controlled trial. *Br J Psychiatry.* 1999;174(1):56-62.
  33. Moore AM, Crane D. Relational diagnosis and psychotherapy treatment cost effectiveness. *Contemporary Family Therapy: An International Journal.* 2014;36(2):281-299.
  34. Flores G, Bridon C, Torres S, et al. Improving asthma outcomes in minority children: a randomized, controlled trial of parent mentors. *Pediatrics.* 2009;124(6):1522-1532.
  35. Dydyk BJ, French G, Gertsman C, Morrison N, O'Neill I. Admitting whole families: an alternative to residential care. *Can J Psychiatry.* 1989;34(7):694-699.
  36. EuroQol Group. EuroQola new facility for the measurement of health-related quality of life. *Health Policy.* 1990;16(3):199-208.
  37. Pote H, Stratton P, Cottrell D, Shapiro D, Boston P. Systemic family therapy can be manualized: Research process and findings. *J Fam Ther.* 2003;25(3):236-262.
  38. Drummond M, Sculpher MJ, Torrance GW, et al. *Methods for the economic evaluation of health care programmes.* 2<sup>nd</sup> ed. Oxford, UK: Oxford Medical Publications, 1997.
  39. Oluboyede Y, Tubeuf S, McCabe C. Measuring health outcomes of adolescents: report from a pilot study. *Eur J Health Econ.* 2013;14(1):11-19.
  40. Adlard N, Kinghorn P, Frew E. Is the UK NICE reference case influencing the practice of pediatric quality-adjusted life-year measurement within economic evaluations? *Value Health.* 2014;17(4):454-461.
  41. Thorrington D, Eames K. Measuring health utilities in children and adolescents: a systematic review of the literature. *PLoS ONE.* 2015;10(8):e0135672.
  42. Griebsch I, Coast J, Brown J. Quality-adjusted life-years lack quality in pediatric care: a critical review of published cost-utility studies in child health. *Pediatrics.* 2005;115(5):e600-e614.
  43. Crane DR. Does family therapy reduce health care costs for more than the identified patient? *Clin Child Psychol Psychiatry.* 2011;16(1):3-4.
  44. Cottrell D, Boston P. Practitioner review: The effectiveness of systemic family therapy for children and adolescents. *J Child Psychol Psychiatry.* 2002;43(5):573-586.
  45. Carr EG. Evidence-based practice in family therapy and systemic consultation I: child focused problems. *J Fam Ther.* 2000; 22(1):29-60.
  46. Asen E. Outcome research in family therapy. *Adv Psychiatr Treat.* 2002;8(3):230-238.